

Research article

Open Access

CD8 α is expressed by human monocytes and enhances Fc γ R-dependent responses

Derrick J Gibbings, Marcelo Marcet-Palacios, Yokananth Sekar, Marcus CY Ng and A Dean Befus*

Address: Pulmonary Research Group, Division of Pulmonary Medicine, Department of Medicine, University of Alberta, Canada

Email: Derrick J Gibbings - gibbings@ualberta.ca; Marcelo Marcet-Palacios - marcelo@ualberta.ca; Yokananth Sekar - yokananth@ualberta.ca; Marcus CY Ng - marcusn@ualberta.ca; A Dean Befus* - dean.befus@ualberta.ca

* Corresponding author

Published: 1 August 2007

Received: 26 February 2007

BMC Immunology 2007, 8:12 doi:10.1186/1471-2172-8-12

Accepted: 1 August 2007

This article is available from: <http://www.biomedcentral.com/1471-2172/8/12>

© 2007 Gibbings et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: CD8 α enhances the responses of antigen-specific CTL activated through TCR through binding MHC class I, favoring lipid raft partitioning of TCR, and inducing intracellular signaling. CD8 α is also found on dendritic cells and rat macrophages, but whether CD8 α enhances responses of a partner receptor, like TCR, to activate these cells is not known. TCR and FcR, use analogous or occasionally interchangeable signaling mechanisms suggesting the possibility that CD8 α co-activates FcR responses. Interestingly, CD8 α + monocytes are often associated with rat models of disease involving immune-complex deposition and FcR-mediated pathology, such as arthritis, glomerulonephritis, ischaemia, and tumors. While rat macrophages have been shown to express CD8 α evidence for CD8 α expression by mouse or human monocytes or macrophages was incomplete.

Results: We detected CD8 α , but not CD8 β on human monocytes and the monocytic cell line THP-1 by flow cytometry. Reactivity of anti-CD8 α mAb with monocytes is at least partly independent of FcR as anti-CD8 α mAb detect CD8 α by western blot and inhibit binding of MHC class I tetramers. CD8 α mRNA is also found in monocytes and THP-1 suggesting CD8 α is synthesized by monocytes and not acquired from other CD8 α + cell types. Interestingly, CD8 α from monocytes and blood T cells presented distinguishable patterns by 2-D electrophoresis. Anti-CD8 α mAb alone did not activate monocyte TNF release. In comparison, TNF release by human monocytes stimulated in a FcR-dependent manner with immune-complexes was enhanced by inclusion of anti-CD8 α mAb in immune-complexes.

Conclusion: Human monocytes express CD8 α . Co-engagement of CD8 α and FcR enhances monocyte TNF release, suggesting FcR may be a novel partner receptor for CD8 α on innate immune cells.

Background

CD8 α is a surface glycoprotein typically found on a sub-population of CTL [1]. CD8 α enhances responses insti-

gated through the TCR by binding MHC class I and signaling through the src kinase lck and the adaptor protein Linker for Activation of T cells (LAT) [2]. The classical

co-receptor model of CD8 suggests CD8 enhances CTL activation by binding the same MHC class I-peptide as TCR [3]. Other evidence suggests CD8 is recruited to the site of T cell activation [4,5]. and can enhance T cell responses even when it does not bind at detectable levels to the same MHC class I-peptide as TCR (e.g. CD8 enhances activation of T cells with an MHC class II specific TCR [6,7]).

CD8 on T cells co-activates responses initiated by TCR, but no such co-activating role has been described for CD8 on other CD8+ cells like dendritic cells [8], NK cells [9,10], mast cells [11] or macrophages (M ϕ) [12]. Interestingly, the Fc γ chain, a component of several FcR [13], NK receptors [14], and ILT1 [15] can substitute for CD3 ζ in TCR expression [16,17]; signaling [18] and T cell activation [19,20] Reciprocally, CD3 ζ can substitute for Fc γ in FcR signaling [21]. Fc γ chain is an ancestral homologue of the CD3 ζ chain [22]. Furthermore, CD3 ζ ^{-/-} η ^{-/-} mice use Fc γ in TCR signaling and CD8-dependent CTL cytotoxicity [19], strongly suggesting CD8 can function with Fc γ in the absence of CD3 ζ or η . In fact, human but not mouse mature T cells often express Syk and Fc γ alongside ZAP-70 and CD3 ζ and in at least some mature effector T cells Syk and Fc γ replace ZAP-70 and CD3 ζ in TCR signaling [23,24]

The cell types that express CD8 α differ among mice, rats and humans. While human [9] and rat NK cells express CD8 α , mouse NK cells do not [25]. Rat M ϕ express CD8 α [12], however, our efforts and those of others to detect CD8 α protein on mouse monocytes and M ϕ have been unsuccessful [26,27]. A portion of CD8 α and all the CD8 β found on mouse dendritic cells is derived from T cells [28]. As transfer of transmembrane proteins between cells is frequently detected, like CD8 in the case above, it is necessary to determine the source and functionality of CD8 α when it is detected on a new cell type or in a new species. Since this study was started, two studies identified binding of anti-CD8 α mAb at high levels to a small percentage of human monocytes during immune responses [29,30] Unfortunately neither study queried whether lower levels of CD8 α were constitutively found on monocytes, demonstrated the cellular origin of the CD8 α found on monocytes, or demonstrated a function for CD8 α on monocytes.

In this report, we provide evidence that human monocytes express CD8 α and that CD8 α can enhance responses mediated through FcR.

Results

CD8 α and not CD8 β is present on human peripheral blood monocytes

Performing flow cytometry on PBMC, a subpopulation of lymphocytes (FSC/SSC gated) expressed high levels of CD8 α and CD8 β , as expected (Figure 1B, anti-CD8 α mAb OKT8 and Figure 1C, anti-CD8 β -dependent mAb 2ST8.5H7). Six anti-CD8 α mAb also bound monocytes at levels greater than three times the geometric mean of isotype mAb (Figure 1B, gated for analysis by expression of high levels of CD14 [31] and characteristic FSC/SSC scatter [Figure 1A]). The monocytic cell line THP-1 bound CD8 α mAb at levels comparable to blood monocytes (data not shown). CD8 β was not detected on monocytes with mAb 2ST8.5H7 (Figure 1C) or 5F2 (not shown), suggesting they do not express CD8 $\alpha\beta$ (mAb 2ST8.5H7) or putative CD8 $\beta\beta$ dimers (mAb 5F2) [32]. Accordingly, mRNA for CD8 β was detected in total PBMC containing CD8 $\alpha\beta$ + T cells, but not in highly enriched monocytes (data not shown).

CD64 does not contribute to anti-CD8 α mAb binding to monocytes

To examine whether Fc receptors contributed to binding of anti-CD8 α mAb to monocytes we first tested the contribution of CD64, the high affinity FcR. CD64 binds immunoglobulin with 100-fold or more the affinity of other Fc γ R [33], is the only Fc γ R that binds monomeric Ig [34] and preferentially binds mouse IgG2a antibodies compared to mouse IgG1 [35]. Accordingly, as human monocytes express significant amounts of CD64 [31] mouse IgG2a has a 100–1000 fold higher affinity for binding human monocytes than mouse IgG1 [36]. Incubating monocytes with human Ig preparations [37] (not shown) or a mAb which blocks binding of Ig to CD64 (clone 10.1) [38] slightly decreased binding of IgG2a isotype control mAb (11–27% geometric mean) and did not decrease binding of anti-CD8 α mAb (Figure 2A).

CD8 α is detected in monocytic cell line and >99% human monocytes independent of FcR

To test whether CD8 α is detected in human monocytes by a method generally acknowledged to be independent of Fc γ R binding to IgG we performed western blot for CD8 α . Proteins at 32 kDa, consistent with CD8 α were detected with anti-CD8 α mAb D9 by western blot (Figure 2B) of thymus lysate, blood lymphocytes, immature monocytes (THP-1), mature *ex vivo* monocytes (>99% CD14^{hi}, CD3 ζ [-ve], enrichment Figure 4A), and M ϕ differentiated with GM-CSF from blood monocytes, but not in lung epithelial cells (A549, negative control). Similarly, a 32 kDa protein was found by western blot with anti-CD8 α mAb B9.11 in PBMC and THP-1 (Figure 2B). These data suggest that anti-CD8 α mAb binding to monocytes is due to the




Figure 1

CD8α is detected by flow cytometry on CD14^{hi} monocytes from human peripheral blood. Anti-CD8α mAb bind monocytes by flow cytometry. **A.** Monocytes were gated by characteristic FSC/SSC scatter (left panel) and expression of high levels of CD14 (right panel) for all flow cytometry studies in this article. **B.** A population of lymphocytes, gated by characteristic FSC/SSC scatter, bind high levels of anti-CD8α. Blood monocytes gated in **A.** bind anti-CD8α mAb. Geometric means of monoclonal antibody binding are shown. Results are representative of five experiments. **C.** Anti-CD8β mAb bind lymphocytes but not monocytes.

presence of CD8α protein and not non-specific binding to FcγR.

Peripheral localization of CD8α on human monocytes: confocal microscopy

To confirm the expected localization of CD8α to the cell periphery on monocytes we performed two-color confocal microscopy of permeabilized PBMC (Figure 3). All anti-CD8α mAb detected CD8α at the periphery of CD14^{hi} monocytes and some CD3^{hi} T cells (Figure 3B–E mAb B9.11 is shown, and is representative of results obtained with LT8, OKT8, 32-M4, 51.1, and Nu-Ts/c). CD8α was also observed intracellularly in some monocytes with a distribution resembling CD14, suggesting that similar to NK cells [39] a small proportion of CD8α

may be found intracellularly in monocytes, perhaps in recycling endosomes.

Human monocytes express CD8α mRNA

The monocytic cell line THP-1 must synthesize CD8α as no other sources of CD8α are available (FBS was CD8α negative by parallel western blot). In contrast *ex vivo* monocytes may acquire CD8α from other CD8α⁺ cells in the body. The presence of CD8α mRNA would suggest that monocytes can synthesize the CD8α protein associated with them. Due to the sensitivity of RT-PCR for mRNA from contaminating cells, we studied the cultured monocytic line THP-1 in addition to highly enriched monocytes (negative for T cell/NK cell specific CD3ζ mRNA, >99% FSC/SSC and CD14^{hi} monocytes, Figure




Figure 2

Binding of anti-CD8α mAb to monocytes is independent of FcR. **A**, Blockade of Ig binding to CD64 with anti-CD64 mAb decreases binding of isotype mAb but not anti-CD8α mAb to monocytes. Bracketed numbers are geometric means of indicated peaks. **B**, Western blot with anti-CD8α D9 detects a 32 kDa protein as expected for CD8α in THP-1, peripheral blood lymphocytes, thymus lysate, peripheral blood monocytes (>99%), GM-CSF differentiated Mφ, PBMC, thymus, and a CTL clone but not in the lung epithelial line A549 (CD8α negative control). Right, anti-CD8α mAb B9.11 detects a 32 kDa protein as expected for CD8α in THP-1 and peripheral blood lymphocytes. 1–1.5 × 10⁶ cell equivalents were loaded in each lane.

4A). CD8α mRNA was detected in peripheral blood mononuclear cells (containing CD8α+ T cells [positive control]), THP-1 monocytic cell line, and highly purified blood monocytes, but not in a lung epithelial cell line (A549) as expected (Figure 4B).

CD8α on monocytes binds MHC class I

Whatever the eventual cellular derivation of CD8α protein found on *ex vivo* monocytes, its ability to function (e.g. bind MHC class I) and impact monocyte responses is practically relevant. We tested whether CD8α on human monocytes contributes to monocyte binding to MHC class I. We expected that anti-CD8α mAb would not block all tetramer binding to human monocytes because members of the immunoglobulin-like-transcript (ILT/CD85)

family (ILT2, 4) expressed by monocytes bind MHC class I tetramers, interact with regions on MHC class I that overlap with CD8α and thus compete with CD8α for binding of MHC class I [40].

Thymocytes from CD8β knockout mice bind MHC class I tetramers and overexpression of CD8α enhances this CD8-dependent binding [41], suggesting that despite the heightened ability of CD8αβ (at least in an unsialylated form on thymocytes [42,43]) to bind MHC class I tetramers, CD8αα is also capable of mediating tetramer binding to T cells.

HLA-*0201 tetramers bound to nearly all CD14^{hi} monocytes (Fig 5A). Tetramers complexed with two different




Figure 3

CD8 α is detected by confocal microscopy on peripheral blood monocytes and lymphocytes with several anti-CD8 α mAb. A, CD3-FITC and CD14-FITC binding to PBMC (Green). B-E, Anti-CD8 α mAb (D, E) binding to monocytes and lymphocytes in comparison to isotype mAb (B, C) (Red). Results are representative of other anti-CD8 α mAb (OKT8, 51.1, 32-M4, Nu-Ts/c, and B9.11).




Figure 4

Human monocytes express CD8α mRNA. Right, purification of monocytes (>99% CD14^{hi} after sorting) was also confirmed by attempting to detect the T and NK cell transcript, CD3ζ, in monocytes. A 293 bp fragment of CD3ζ mRNA was detected in PBMC, containing T cells, but not monocytes, the monocytic line THP-1, or lung epithelial cells (A549) after 50 cycles. B, A 379 bp CD8α mRNA fragment was detected by RT-PCR in THP-1, >99% CD14^{hi} monocytes, PBMC, but not in a lung epithelial cell line, (A549) using intron-spanning primers, and 35 cycles of cDNA amplification. Detection of β-actin mRNA confirmed RNA extraction and RT-PCR was performed successfully.

peptides bound similarly to monocytes (data not shown). In agreement with others [44,45], we found that anti-CD8α clone B9.11 inhibited HLA tetramer binding (Figure 5B, 11.6%). Another anti-CD8α clone, D9, also inhibited tetramer binding (Figure 5B, 18.6%, $p < 0.05$). Finally, tetramer binding was not affected by clone 32-M4 (despite its ability to bind CD8α on monocytes), or isotype control mAb (Figure 5B). Other studies of MHC class I tetramer binding to CD8α using several anti-CD8α mAb have also shown that tetramer binding may be unaffected, inhibited or enhanced by anti-CD8α mAb, depending on which anti-CD8α clone is used, and TCR affinity [43,46,47].

While our evidence suggests CD8α contributes to MHC class I tetramer binding by human monocytes, the

observed diminution of tetramer binding may be via an effect on the kinetics or stability of tetramer binding to other receptors for MHC class I.

Fewer Sialylated 34 kDa Versions of CD8α are found on Monocytes Compared to T cells

Previous publications have demonstrated notable differences in immunoprecipitated CD8α by 2-D electrophoresis. Some authors detected immunoprecipitated CD8α from thymus as restricted spots of pI ~ 6 [48]. In contrast, others detected immunoprecipitated CD8α from blood at numerous spots ranging from pI 6–9.5 [49], and molecular weights of 32 to 34 kDa [50]. We tested whether CD8α from monocytes in comparison to blood lymphocytes exhibit a distinct pattern of sialylation or other post-translational pattern detectable by 2-D electrophoresis.




Figure 5

CD8 α on human peripheral blood monocytes mediates MHC class I binding. **A**, Dark histogram is background fluorescence of CD14^{hi} monocytes, light histogram represents binding of PE-labeled tetramers. **B**, Bar graph is pooled results of gated CD14^{hi} monocytes from three different donors in separate experiments. Inhibition of tetramer binding is expressed as percent decrease in mean fluorescence intensity of tetramer binding due to pretreatment with anti-CD8 α mAb compared to isotype mAb (* $p < 0.05$, non-paired t-test).

A polyclonal anti-CD8 α antibody detected discrete spots across 2-D gels from the predicted pI of unglycosylated CD8 α (~9) to pI 6–7, at Mr from 32–34 kDa (Figure 5), as shown by others [49]. Much of the heterogeneity detected with anti-CD8 α polyclonal antibody could be eliminated by treatment with neuraminidase, suggesting it is due to sialylation of CD8 α (Figure 6A). Notably, monocytes had less of the sialylated 34 kDa forms of CD8 α than T cells in samples from three individuals (Figure 6A). In contrast, anti-CD8 α clone D9 detected a single protein spot (32 kDa, ~pI 6), in a pattern similar to that found by others [48]. To confirm that the protein recognized at 32 kDa pI 6 was CD8 α we sequenced it by MALDI-QTOF from 2-D gels (see Additional file 1). Neuraminidase treatment

diminished but did not eliminate the protein spot recognized by anti-CD8 α clone D9 and gave rise to faint basic spots similar to those observed with polyclonal anti-CD8 α antibody after neuraminidase treatment. D9 may preferentially recognize particular glycosylation/sialylation patterns of CD8 α (Figure 6B,C).

In our experiments large differences in 2-D electrophoresis patterns for CD8 α (e.g. between D9 and polyclonal anti-CD8 α Ab) are due to the specificity of different antibodies recognizing CD8 α , and not a result of differences in the cell type expressing CD8 α . Nonetheless, using the same polyclonal antibody, subtle differences in CD8 α were




Figure 6

Subtle differences in CD8α between monocytes and lymphocytes detected by 2-D electrophoresis. A, Western blot with a polyclonal anti-CD8α antibody after 2-D electrophoresis of *ex vivo* monocytes and lymphocytes. Cell lysates of lymphocytes and monocytes from one donor were separated by adherence, halved and treated (bottom panels) or not treated (top panels) with neuraminidase before analysis. B, Western blot with anti-CD8α mAb D9. C, Neuraminidase treatment and western blot with D9 of lymphocyte lysate. Results are representative of experiments with three donors.

found between monocytes and lymphocytes by 2-D electrophoresis.

Anti-CD8α mAb amplifies monocyte responses to immune-complexes through FcγR

TCR and FcR use analogous and sometimes interchangeable signaling mechanisms to activate cells [20], and CD3ζη-null mice can use Fcγ to reconstitute CD8/TCR-

dependent CTL cytotoxicity[23]. We investigated if CD8α on monocytes might be involved in responses to immune-complexes mediated by FcγR. To this end, we used a common immune-complex-FcγR activation system.

Treatment of monocytes with monomeric non-specific IgG_{2a} mAb induced a slight increase in monocyte TNF pro-

duction (Figure 7A,B). This is likely due to the ability of the high affinity Fc γ R, CD64, found on the majority of monocytes to bind monomeric mouse IgG_{2a} [36]. Monomeric anti-CD8 α mAb did not increase monocyte TNF production more than control IgG_{2a} mAb (Figure 7A,B). When monocytes were stimulated with preformed immune-complexes (isotype mAb cross-linked with anti-mouse Ig), CD14^{hi} monocytes produced moderate amounts of intracellular TNF (Figure 7A,B). Formation of immune-complexes with anti-CD8 α mAb (32-M4 cross-linked with anti-mouse Ig) rather than isotype mAb, resulted in production of 2-fold more TNF by monocytes (Figure 7A,B). Immune-complexes containing anti-CD8 α mAb D9 did not significantly increase monocyte TNF production above control immune-complexes, indicating that monocyte TNF release is not significantly stimulated by any immune-complex containing a mAb that binds to the monocyte surface. It is not surprising that only one of two anti-CD8 α mAb enhanced Fc γ R-dependent responses, because when others have screened several anti-CD8 α mAb in parallel for effects on CTL cytotoxicity, or CD8-MHC class I binding (see above), the effect ranged from substantial inhibition to no effect depending on the particular anti-CD8 α mAb clone [51-53]

Enhanced activation of monocytes incubated with immune-complexes containing anti-CD8 α mAb in comparison to immune-complexes with isotype control was also found when TNF release (1376 pg/mL versus 468 pg/mL, $p < 0.05$ [data not shown]), CD69 upregulation, or CD14 downregulation was measured (Figure 7C).

Blockade of FcR with excess Fc fragments nullified TNF release instigated by isotype and anti-CD8 α immune-complexes (Figure 7), suggesting the immune-complex system used is Fc γ R-dependent, as expected. As CD8 α enhancement of TNF production is inhibited by Fc fragments and does not occur with monomeric anti-CD8 α mAb, the ability of anti-CD8 α mAb to enhance responses of human monocytes appears to depend on co-engagement of FcR.

Discussion

The present study demonstrates expression of CD8 α by monocytic cells, and suggests that CD8 α , in addition to co-activating TCR responses may have a previously unacknowledged role in co-activating Fc γ R responses.

Previous to the present study no evidence existed to demonstrate that human monocytes synthesized CD8 α . Human PBMC can appear CD14⁺/CD8 α ^{hi} in HIV infected individuals, but this population is due to acquisition of CD14 from monocytes, by T cells, that are mostly CD8 α ⁺ [54,55]. A similar effect may explain previous claims of CD8 α on monocytes subsequent to dengue virus infec-

tion [30]. Other studies have demonstrated that macrophages and dendritic cells can acquire CD8 α from T cells, without themselves synthesizing it [28]. Thus, the detection of CD8 α protein on monocytes in other studies [29] does not demonstrate that CD8 α is expressed by, or more importantly, functional on monocytes.

We provide strong evidence that human monocytes constitutively express CD8 α at low levels. Notably, CD8 α was observed in monocytes, and the monocytic cell line THP-1 by western blot confirming the presence of CD8 α in these cells without possible contribution of Fc γ R. Moreover, detection of CD8 α mRNA in THP-1, CD8 α protein on THP-1 and 32 kDa CD8 α in lysate of continuously cultured THP-1, demonstrates that these monocytic cells must transcribe CD8 α mRNA and translate CD8 α protein.

We find that monocytes have less 34 kDa sialylated forms of CD8 α compared to T cells. The Mr difference between 32 and 34 kDa forms of CD8 α may be due to glycosylation other than sialylation [50] (Figure 5), or palmitoylation of CD8 α [56] at three eligible membrane-proximal cysteines [57]. Monocytes and T cells may express different glycosylation or palmitoylation enzymes that account for predominant accumulation of 32 or 34 kDa CD8 α . Phosphorylation of CD8 α , or oxidation of the free cysteine in the CD8 α Ig-domain may explain pI differences of CD8 α remaining after neuraminidase treatment. Sialylation of CD8 α [58], like CD8 β [42], changes during T cell development, and potentially upon T cell activation. Differential sialylation and palmitoylation of CD8 are known to modulate its ability to bind MHC class I and induce T cell activation [42,43,59] Unfortunately, we cannot relate the differences in CD8 α we observe between monocytes and T cells to their comparative ability to bind MHC class I and co-activate signaling.

The ability of CD8 to enhance T cell responses in a TCR-dependent manner was defined over twenty years ago. Since then, CD8 α expression by several cell types that lack the TCR such as human NK cells [9], dendritic cells [60], rat macrophages [12] and mast cells [61] has been established. Some evidence has suggested a role for CD8 α in apoptosis and survival (e.g. [39]) however a viable model has not emerged to define when and how CD8 α activation by its ubiquitous MHC class I ligand may be controlled on these cells that lack the TCR.

MHC class I-like protein TL, which exhibits a restricted expression pattern and a high binding affinity for mouse CD8 α , but not CD8 β [62] could hypothetically control activation of CD8 α ⁺ monocytes or dendritic cells. One model has proposed that mouse CD8 α may downregulate T cell responses and promote generation of memory T cells by binding TL [63,64] however, evidence that




Figure 7

CD8 α enhances monocyte TNF production in a Fc γ R-dependent manner. Monocytes were incubated with anti-CD8 α or isotype mAb alone or in immune-complexes. Excess Fc fragment was used to block Fc γ R. TNF production was measured by intracellular flow cytometry after 5 h (gated on CD14^{hi} monocytes). A. Histograms are representative results. B. Bar graph represents average geometric mean of intracellular TNF from four separate experiments with monocytes from different donors. Standard error of the mean is designated by error bars. *: $p < 0.05$ non-paired t-test. C. CD69 and CD14 regulation induced by immune-complexes (IC) is enhanced by CD8 α ligation (representative example).

CD8 α has a role in generation of memory T cells is controversial [65-67]. Similarly, while one study suggests an inhibitory role of CD8 α on proliferation and cytotoxicity of T cells (but an enhancement of cytokine release) [68], several studies demonstrate the ability of CD8 α to enhance T cell cytotoxicity and other responses [69-73]. In the absence of a equivalent of TL in humans that binds CD8 α with high affinity, the applicability of this model to humans is even more problematic.

Interestingly, some evidence previous to this report suggested CD8 may be able to co-activate responses of Fc γ R. Many of the components of FcR and TCR signaling are homologous or interchangeable, such as Fc γ and CD3 ζ chain, ZAP-70 and Syk [74,75], or LAT [76]. For instance, Fc γ and CD3 ζ are conserved ancestral duplicates [77] that can substitute for each other in activation of mast cells, $\gamma\delta$ or $\alpha\beta$ T cells [18,20,23] LAT binds CD8 α , and is phosphorylated upon macrophage activation through Fc γ R [76] or upon Fc γ -dependent activation in mast cells and platelets [78]. A more direct suggestion that CD8 can co-activate Fc γ -chain dependent signaling through TCR was provided by the demonstration of CD8 involvement in TCR-mediated cytotoxicity in Fc γ +, CD3 $\zeta\eta$ -/- CTL [19]. What is more, Fc γ and Syk are naturally expressed by many mature and immature T cells in humans (less so in mice), and participate in TCR-signaling [23], suggesting that *in vivo* even on T cells CD8 may have a role in activating Fc γ responses[24]. In sum, reasonable although little acknowledged evidence existed that CD8 participated in Fc γ chain-linked responses in T cells. However, no evidence had shown whether CD8 could enhance Fc γ chain responses that were not mediated through TCR thereby expanding this model to include possible functions of CD8 on monocytes, NK cells, dendritic cells, or mast cells.

We and others have previously demonstrated that CD8 α on rat M ϕ and NK cells signals through Syk and src tyrosine kinases [79], consistent with TCR and Fc γ R signaling, but in the absence of TCR [80,81] Here, we find that anti-CD8 α mAb enhances TNF production of monocytes exposed to immune complexes in an Fc-dependent manner, mirroring the ability of CD8 α to enhance T cell activation in a TCR-dependent manner. This evidence suggests CD8 α enhances Fc γ R responses, through at least one of several potential mechanisms. In experiments presented here CD8 α may promote initial contact and binding stability of anti-CD8 α mAb containing immune-complexes with Fc γ R. In the same way, CD8 α may promote binding of Fc γ R to MHC class I-expressing cells coated with immune-complexes in cancer, viral infection or autoimmune disease. Alternatively, or in addition, signaling through CD8 α may enhance activation of Fc γ R signaling in our experiments. While we have not directly demonstrated that CD8 α signaling enhances Fc γ R signal-

ing, previous evidence supports this possibility and suggests it merits further investigation. If CD8 α signaling can enhance Fc γ R signaling then hypothetically CD8 α may enhance responses of other receptors that both CD3 ζ and Fc γ can function with such as NKp30, and NKp46 [14] in NK cells, and Fc ϵ RI in mast cells [21].

Our evidence suggests monocyte responses instigated through immune-complexes and FcR can be amplified by co-engagement of CD8 α . Interestingly, in rats CD8 α + monocytes and M ϕ are found at sites of tissue damage in immune-complex mediated glomerulonephritis [82], arthritis [83], tumor [84], experimental allergic encephalomyelitis (a model of multiple sclerosis) [85], and ischaemia-reperfusion injury [86]. TNF is an important mediator in many of these diseases [87,88] As monocyte TNF production is enhanced by co-activation of CD8 α and immune-complexes, CD8 α on monocytes may aggravate some autoimmune and acute inflammatory conditions characterized by tissue deposition of immune-complexes.

In summary, we find that human monocytes express CD8 α and that monocyte CD8 α is differentiable from that on T cells by 2-D electrophoresis. We provide evidence that CD8 α on monocytes amplifies responses initiated through FcR, suggesting for the first time a co-activator role for CD8 on cells without the TCR.

Methods

Antibodies

Isotype control antibodies were mouse IgG₁ and IgG_{2a} (Sigma, St. Louis, MO), IgG_{2a}-FITC, and -PE, (Caltag, Burlingame, CA). Anti-CD8 α mAb used were: D9 and 32-M4 (Santa Cruz, Santa Cruz, CA) LT8 (Serotec, Raleigh, NC), B9.11 (Beckman-Coulter Canada Inc., Mississauga, ON), and Nu-Ts/c (Nichei Corp., Tokyo, Japan). Polyclonal anti-human CD8 α Ab (H160) was obtained from Santa Cruz. Anti-CD8 α mAb 51.1 (gift of Dr. D. Burshtyn, University of Alberta) and OKT8, and anti-rat MHC class I mAb OX18 (European Collection of Cell Cultures, Salisbury, UK) were purified from hybridoma supernatant by protein G affinity chromatography. Anti-CD8 β -antibodies were obtained from Beckman-ImmunoTech (clone 2ST8.5H7-PE, Mississauga, Canada) and Serotec (clone 5F2). Anti-CD3-FITC and anti-CD14-FITC/PE were obtained from Caltag. Anti-mouse Ig-FITC (STAR70) was obtained from Serotec. Anti-CD69 mAb and matching isotype control were obtained from BD Biosciences. Anti-mouse Ig-HRP was purchased from Pierce (Rockford, IL).

Cell recovery and culture

The promonocytic cell line THP-1 was maintained in American Type Culture Collection recommended media (RPMI 1640 medium, 2 mM L-glutamine, 1.5 g/L sodium

bicarbonate, 4.5 g/L glucose, 10 mM HEPES, 1.0 mM sodium pyruvate 0.05 mM 2-mercaptoethanol, and 10% fetal bovine serum [FBS]). CTL clones [68] were a gift of Dr. Chris Bleackley (University of Alberta).

Human blood (100 mL) was collected into heparinized tubes. Red blood cells were sedimented by addition of 7 mL 6% dextran (Sigma) in RPMI 1640 per 35 mL blood (0.5 h, room temperature). PBMC were enriched on Ficoll-Paque Plus (Amersham Biosciences, Oakville, ON, Canada) and washed three times in PBS. Monocytes were further enriched by three methods. Greater than 80% enriched monocytes were obtained from a Percoll gradient [89] for studies of monocyte activation. Monocytes enriched by Percoll were further purified (>99%) by anti-CD14-PE flow sorting for western blot and RT-PCR analysis. M ϕ were differentiated from adherent monocytes (1 h, 37 C) with 500 ng/mL GM-CSF (Biosource, Camarillo, CA) for 3 d.

Flow cytometry

Cells on ice were blocked with 5% milk, 0.1% bovine serum albumin (BSA) in PBS. In some experiments human Ig (50 μ g/mL, Bethyl Laboratories Inc., Montgomery, TX) was used to minimize binding of mAb to FcR. Cells were treated with 10 μ g/mL isotype mAb or anti-CD8 α mAb, washed three times and incubated with anti-mouse Ig-FITC Ab (1/100, STAR70, Serotec). Anti-CD8 β mAb 2ST8.5H7 directly conjugated to PE was used at 10 μ g/mL and compared to IgG2a-PE. Cells were washed three times and incubated with 1/10 normal mouse serum before addition of anti-CD14-FITC (1/50).

To analyse the contribution of high affinity Fc γ RI to anti-CD8 α mAb binding to monocytes, PBMC were pre-incubated for 30 min with anti-CD64 mAb clone 10.1 (10 μ g/mL, BioLegend, San Diego, CA), which blocks binding of Ig to CD64 [38]. Cells were washed and incubated with IgG2a-PE (10 μ g/mL, BD Biosciences) or 32-M4-PE (10 μ g/mL, Santa Cruz). Cells were washed and data was collected on a FACScan.

All flow cytometry analysis was performed with WinMDI and CellQuest Pro (BD Biosciences) programs. Monocytes were gated by characteristic FSC/SSC scatter and high expression of CD14.

Confocal Microscopy

PBMC were adhered to poly-L-lysine coated coverslips for 0.5 h, fixed with 4% paraformaldehyde (10 min) and permeabilized with 0.1% triton-X-100 in PBS (10 min). Cells were blocked (10% FBS, 3% BSA, 30 min) before staining with anti-CD8 α mAb (10 μ g/mL). Cells were washed three times (5 min, 2 mL PBS) between each reagent. Cells were sequentially incubated with anti-mouse-Ig-rhodam-

ine red (Molecular Probes, Eugene, OR), 1/10 normal mouse serum and anti-CD14-FITC or anti-CD3-FITC (Caltag). Images were obtained using an Olympus FV1000 confocal microscope (Carsen Group, Markham, ON) with Fluoview software.

Western Blot

Cells were lysed in 1% Triton X-100 for 20 min. Lysate was centrifuged for 5 min at 1000 g to eliminate non-solubilized material. Before loading on gels remaining lysate was diluted 2-fold with Laemmli buffer (BioRad), 2% 2-mercaptoethanol was added and samples were boiled for 5 min. Similar amounts of cell lysate (1–1.5 \times 10⁶ cell equivalents) were loaded per lane on 4–20% SDS-PAGE denaturing gels (Bio-Rad Readygel). Wet protein transfers to PVDF were performed at 100 V for 1 h. PVDF was blocked for 1 h with 5% milk in TBS, 0.1% Tween-20 and subsequently blotted with 0.2 μ g/mL primary antibody.

RT-PCR

Total RNA was isolated using the Qiagen RNeasy kit. RNA was reverse-transcribed with Superscript II reverse transcriptase (Invitrogen, Carlsbad, CA) using oligo(dT) as primers. The cDNA concentration was estimated by absorbance for each sample and diluted to 100 ng/25 μ l of reaction volume. The number of cycles was optimized to be in the exponential phase of the reaction by performing the reaction at different cycles. Densitometric analysis of the gels was performed to select optimal PCR cycle numbers. Thereafter, PCR was performed in 20 μ l reactions with primer pairs (25 μ M) below. Intron-spanning primers of the sequence 5'-TTTCGGCGAGATACGCTAACCCTGTGC-3' and 5'-TTTAGCCTC-CCCCTTTGTAACCGGGCG-3' were used to generate a CD8 α cDNA fragment of 379 bp [70]. Intron-spanning primers generating a 209 bp product for CD8 β were 5'-GGTGAAGAGGTGGAACAGGA-3' and 5'-CTTGAGGGT-GGACTTCTTGG-3'. A β -actin cDNA fragment of 326 bp was produced using intron-spanning primers of sequence 5'-GGC ATC CTC ACC CTG AAG TA-3' and 5'-AGG GCA TAC CCC TCG TAG AT-3'. PCR amplification was performed for 35 cycles of 1 min at 94 C, 1 min at 60 C and 2 min at 72 C, and a final cycle of 72 C for 10 min to complete polymerization. PCR products were run on a 1.5% agarose gel containing ethidium bromide. Intron-spanning primers were used and samples were treated with 84 U/ μ l DNase I before RT-PCR to avoid interference of contaminating DNA in purified RNA.

Samples used to amplify CD8 α mRNA were also amplified with intron-spanning CD3 ζ RT-PCR primers (5'-GCACAGTTGCCGATTACAGA-3' and 5'-GCCACGTCTCTGTCCAAA-3', 293 base pair product) for 50 cycles, performed as above.

2-D electrophoresis

Lymphocytes and monocytes were enriched by collecting non-adherent and adherent cells respectively after 1 h in culture flasks. Lymphocyte and monocyte lysates were prepared using the 2-D cleanup kit (Bio-Rad) and resuspended in IPG strip rehydration buffer (Bio-Rad) with 2% carrier isoelectric point (pI) 3–10 ampholytes (Bio-Rad). Lysates were focused on 7 cm pI 3–10 strips (Bio-Rad).

Monoclonal antibody affinity chromatography

OKT8 at 5–10 mg/mL in 0.1 M HEPES pH 7.5 was coupled to pre-washed N-hydroxysuccinimidyl-activated agarose beads (Sigma) at 4 C for 1 h. Remaining active sites were blocked by incubating in the presence of 0.1 mL 1 M ethanolamine pH 8 at 4 C for 1 h. Rat cultured mast cell line (RCMC, MHC class I purification) or human thymus (CD8 purification) was lysed with 1% triton X-100 in PBS with Complete Mini anti-protease cocktail tablets (Roche Applied Science, Laval, PQ, Canada). Supernatant remaining after 1000 g, 12,000 g, and 100,000 g centrifugations was loaded on columns. Columns were sequentially washed with 30 volumes lysis buffer, 20 volumes 10 mM Tris 0.5% triton X-100 300 mM NaCl pH 8, 20 volumes 10 mM sodium phosphate 0.5% triton X-100 450 mM NaCl pH 10, and eluted with 0.05 M diethylamine 0.5% triton X-100 650 mM NaCl pH 11.5. 1.5 mL fractions were collected into 50 µL 1 M Tris HCl pH 6.7.

MALDI-QTOF

Bands were excised and an automated in-gel tryptic digestion was performed on a Mass Prep Station (Water, USA). The gel pieces were de-stained, reduced (DTT), alkylated (iodoacetamide), digested with trypsin (Promega Sequencing Grade Modified) and the resulting peptides extracted from the gel and analyzed via LC/MS/MS. LC/MS/MS was performed on a CapLC HPLC (Waters, USA) coupled with a Q-ToF-2 mass spectrometer (Waters, USA). Tryptic peptides were separated using a linear water/acetonitrile gradient (0.2% Formic acid) on a Picofrit reversed-phase capillary column, (5 micron BioBasic C18, 300 Angstrom pore size, 75 micron ID × 10 cm, 15 micron tip) (New Objectives, MA, USA), with an in-line PepMap column (C18, 300 micron ID × 5 mm), (LC Packings, CA, USA) used as a loading/desalting column. Protein identification from the generated MS/MS data was done searching the NCBI non-redundant database using Mascot Daemon (Matrix Science, UK). Search parameters included carbamidomethylation of cysteine, possible oxidation of methionine and one missed cleavage per peptide.

MHC class I binding

PE labeled HLA-A*0201 tetramers assembled [71] with two peptides selected by the SYFPEITHI search engine, *Mycobacterium tuberculosis* antigen 85-B 143–152 (KLVAN-

NTRL) and a diabetes-specific epitope of glutamic acid decarboxylase 114–123 (VMNILLQYVV) were a gift of Dr. John Elliott (University of Alberta). All washes and incubations of cells were done in ice cold PBS with 0.02% NaN₃. 1 × 10⁶ human PBMC were incubated 15 min with 1.5 µg tetramers before the addition of CD14-FITC for 15 min. Before flow cytometry analysis cells were washed three times. In some experiments cells were incubated with 40 µg/mL anti-CD8α mAb or isotype control for 30 min prior to addition of tetramers.

Immune-Complex Stimulation of Monocytes and TNF Measurements

Monocytes were treated with isotype or anti-CD8α mAb (10 µg/mL), or immune-complexes. All mAb were negative for endotoxin using the LAL assay (Sigma-Aldrich). Immune-complexes were prepared by combining isotype mAb or anti-CD8α mAb (10 µg/mL) with anti-mouse Ig (20 µg/mL) for 15 min before addition to monocytes. Monocytes enriched on a Percoll gradient were incubated with immune-complexes for 5 h at 0.2 × 10⁶ cells/well of a 96 well plate (Becton Dickinson, 35172). Anti-human TNF-PE mAb, fixation and permeabilization buffers, and monensin were from ebioscience (San Diego, CA). Intracellular TNF was detected according to supplier recommendations. Monensin (2 µM) was added 2 h after immune-complexes.

Stimulation of monocytes with immune-complexes was inhibited by pretreating monocytes with 50 µg/mL purified mouse IgG Fc fragment (Jackson Immunoresearch, West Grove, USA) to block binding of immune-complexes to FcR. In these experiments free binding sites of anti-mouse Ig antibody in immune-complexes that might otherwise bind Fc fragments on pretreated monocytes, were pre-blocked with 50 µg/mL purified Fc fragment.

TNF release was measured by ELISA after activation of monocytes for 18 h using 2 µg/mL anti-human TNF mAb as a capture antibody (clone 28401, R&D Systems, Minneapolis, MN) and biotinylated anti-human TNF (BAF210, R&D systems) as a detection antibody. Signal was detected with streptavidin-HRP (Vector Labs, Burlingame, CA) and o-phenylenediamine.

Measurement of CD14 and CD69 Expression

After 18 h activation cells were stained with CD14-FITC/CD69-PE or isotype controls and fixed (ebioscience fixation buffer) before analysis on a FACScan.

Abbreviations

AM (alveolar macrophage), Mφ (macrophage), geometric mean (Gm), immunoglobulin-like-transcript (ILT), isoelectric point (pI), Linker for Activation of T cells (LAT), peripheral blood mononuclear cell (PBMC)

Authors' contributions

DG designed and analyzed experiments, wrote the manuscript and performed flow cytometry, western blots, 2-D electrophoresis, confocal microscopy, MHC binding and monocyte activation assays. MMP helped perform confocal microscopy and designed, performed and analyzed RT-PCR results for CD8 β and CD3 ζ . YS performed 2D blots with polyclonal antibodies and analyzed results. MCYN performed RT-PCR for CD8 α and some flow cytometry and western blot. ADB designed and analyzed experiments, revised and approved this document. All authors read and approved the final manuscript.

Additional material

Additional file 1

Protein recognized by anti-CD8 α mAb D9 is CD8 α . CD8 α was enriched from human thymus lysate by immunoaffinity chromatography with anti-CD8 α mAb OKT8. OKT8-reactive fractions were analyzed by western blot with anti-CD8 α mAb D9 (left), and silver stain (right) after 2-D electrophoresis. Alignment of western blot and silver stain gels allowed extraction of D9-reactive spots from silver stained gels for peptide sequencing by MALDI-QTOF (lower panel).

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2172-8-12-S1.ppt>]

Acknowledgements

Supported by Canadian Institutes of Health Research (CIHR) and Alberta Lung Association operating grants (ADB), an Alberta Lung Association Studentship award to DG and summer studentship award (MN) from the Alberta Heritage Foundation for Medical Research

References

- Ledbetter JA, Evans RL, Lipinski M, Cunningham-Rundles C, Good RA, Herzenberg LA: **Evolutionary conservation of surface molecules that distinguish T lymphocyte helper/inducer and cytotoxic/suppressor subpopulations in mouse and man.** *J Exp Med* 1981, **153**:310-323.
- Bosselut R, Zhang W, Ashe JM, Kopacz JL, Samelson LE, Singer A: **Association of the adaptor molecule LAT with CD4 and CD8 coreceptors identifies a new coreceptor function in T cell receptor signal transduction.** *J Exp Med* 1999, **190**:1517-1526.
- Janeway CA Jr: **The T cell receptor as a multicomponent signalling machine: CD4/CD8 coreceptors and CD45 in T cell activation.** *Annu Rev Immunol* 1992, **10**:645-674.
- Thome M, Duplay P, Guttinger M, Acuto O: **Syk and ZAP-70 mediate recruitment of p56lck/CD4 to the activated T cell receptor/CD3/zeta complex.** *J Exp Med* 1995:1997-2006.
- Thome M, Germain V, DiSanto JP, Acuto O: **The p56lck SH2 domain mediates recruitment of CD8/p56lck to the activated T cell receptor/CD3/zeta complex.** *Eur J Immunol* 1996:2093-2100.
- Miceli MC, von Hoegen P, Parnes JR: **Adhesion versus coreceptor function of CD4 and CD8: role of the cytoplasmic tail in coreceptor activity.** *Proc Natl Acad Sci USA* 1991, **88**:2623-2627.
- Glaichenhaus N, Shastri N, Littman DR, Turner JM: **Requirement for association of p56lck with CD4 in antigen-specific signal transduction in T cells.** *Cell* 1991, **64**:511-520.
- Winkel K, Sotzik F, Vremec D, Cameron PU, Shortman K: **CD4 and CD8 expression by human and mouse thymic dendritic cells.** *Immunol Lett* 1994:93-99.
- Perussia B, Fanning V, Trinchieri G: **A human NK and K cell subset shares with cytotoxic T cells expression of the antigen recognized by antibody OKT8.** *J Immunol* 1983, **131**:223-231.
- Moebius U, Kober G, Griscelli AL, Hercend T, Meuer SC: **Expression of different CD8 isoforms on distinct human lymphocyte subpopulations.** *Eur J Immunol* 1991, **21**:1793-1800.
- Lin TJ, Hirji N, Nohara O, Stenton GR, Gilchrist M, Befus AD: **Most cells express novel CD8 molecules that selectively modulate mediator secretion.** *J Immunol* 1998, **161**:6265-6272.
- Hirji N, Lin TJ, Befus AD: **A novel CD8 molecule expressed by alveolar and peritoneal macrophages stimulates nitric oxide production.** *J Immunol* 1997, **158**:1833-1840.
- Davis RS, Dennis G Jr, Odom MR, Gibson AW, Kimberly RP, Burrows PD, Cooper MD: **Fc receptor homologs: newest members of a remarkably diverse Fc receptor gene family.** *Immunol Rev* 2002, **190**:123-136.
- Moretta A, Bottino C, Vitale M, Pende D, Cantoni C, Mingari MC, Biassoni R, Moretta L: **Activating receptors and coreceptors involved in human natural killer cell-mediated cytotoxicity.** *Annu Rev Immunol* 2001, **19**:197-223.
- Nakajima H, Samaridis J, Angman L, Colonna M: **Human myeloid cells express an activating ILT receptor (ILT1) that associates with Fc receptor gamma-chain.** *J Immunol* 1999:5-8.
- Petersson K, Ivarsson F: **Early TCR alpha beta expression promotes maturation of T cells expressing Fc epsilon RI gamma containing TCR/CD3 complexes.** *J Immunol* 2001, **166**:6616-6624.
- She J, Simpson SJ, Gupta A, Hollander G, Levelt C, Liu CP, Allen D, van Houten N, Wang B, Terhorst C: **CD16-expressing CD8alpha alpha+ T lymphocytes in the intestinal epithelium: possible precursors of Fc gammaR-CD8alpha alpha+ T cells.** *J Immunol* 1997, **158**:4678-4687.
- Qian D, Sperling AI, Lancki DW, Tatsumi Y, Barrett TA, Bluestone JA, Fitch FW: **The gamma chain of the high-affinity receptor for IgE is a major functional subunit of the T-cell antigen receptor complex in gamma delta T lymphocytes.** *Proc Natl Acad Sci USA* 1993, **90**:11875-11879.
- Liu CP, Lin WJ, Huang M, Kappler JW, Marrack P: **Development and function of T cells in T cell antigen receptor/CD3 zeta knockout mice reconstituted with Fc epsilon RI gamma.** *Proc Natl Acad Sci USA* 1997, **94**:616-621.
- Rodewald HR, Arulanandam AR, Koyasu S, Reinherz EL: **The high affinity Fc epsilon receptor gamma subunit (Fc epsilon RI gamma) facilitates T cell receptor expression and antigen/major histocompatibility complex-driven signaling in the absence of CD3 zeta and CD3 eta.** *J Biol Chem* 1991, **266**:15974-15978.
- Howard FD, Rodewald HR, Kinet JP, Reinherz EL: **CD3 zeta subunit can substitute for the gamma subunit of Fc epsilon receptor type I in assembly and functional expression of the high-affinity IgE receptor: evidence for interreceptor complementation.** *Proc Natl Acad Sci USA* 1990, **87**:7015-7019.
- Ra C, Jouvin MH, Blank U, Kinet JP: **A macrophage Fc gamma receptor and the mast cell receptor for IgE share an identical subunit.** *Nature* 1989, **341**:752-754.
- Krishnan S, Warke VG, Nambiar MP, Tsokos GC, Farber DL: **The FcR gamma subunit and Syk kinase replace the CD3 zeta-chain and ZAP-70 kinase in the TCR signaling complex of human effector CD4 T cells.** *J Immunol* 2003, **170**:4189-4195.
- Chu DH, van Oers NS, Malissen M, Harris J, Elder M, Weiss A: **Pre-T cell receptor signals are responsible for the down-regulation of Syk protein tyrosine kinase expression.** *J Immunol* 1999, **163**:2610-2620.
- Kieffer LJ, Bennett JA, Cunningham AC, Gladue RP, McNeish J, Kavathas PB, Hanke JH: **Human CD8 alpha expression in NK cells but not cytotoxic T cells of transgenic mice.** *Int Immunol* 1996, **8**:1617-1626.
- Leon B, Martinez dH, Parrillas V, Vargas HH, Sanchez-Mateos P, Longo N, Lopes-Bravo M, Ardavin C: **Dendritic cell differentiation potential of mouse monocytes: monocytes represent immediate precursors of CD8- and CD8+ splenic dendritic cells.** *Blood* 2004, **103**:2668-2676.
- Nault G: **Expression of CD8 on mouse alveolar macrophages.** In Thesis (MSc) Edmonton, Alberta, University of Alberta; 2003.

28. Vremec D, Pooley J, Hochrein H, Wu L, Shortman K: **CD4 and CD8 expression by dendritic cell subtypes in mouse thymus and spleen.** *J Immunol* 2000, **164**:2978-2986.
29. Baba T, Ishizu A, Iwasaki S, Suzuki A, Tomaru U, Ikeda H, Yoshiki T, Kasahara M: **CD4+/CD8+ macrophages infiltrating at inflammatory sites: a population of monocytes/macrophages with a cytotoxic phenotype.** *Blood* 2006, **107**:2004-2012.
30. Liu CC, Huang KJ, Lin YS, Yeh TM, Liu HS, Lei HY: **Transient CD4/CD8 ratio inversion and aberrant immune activation during dengue virus infection.** *J Med Virol* 2002, **68**:241-252.
31. Grage-Griebenow E, Flad HD, Ernst M: **Heterogeneity of human peripheral blood monocyte subsets.** *J Leukoc Biol* 2001, **69**:11-20.
32. Devine L, Kieffer LJ, Aitken V, Kavathas PB: **Human CD8 beta, but not mouse CD8 beta, can be expressed in the absence of CD8 alpha as a beta beta homodimer.** *J Immunol* 2000:833-838.
33. Ravetch JV, Bolland S: **IgG Fc receptors.** *Annu Rev Immunol* 2001, **19**:275-290.
34. Nimmerjahn F, Ravetch J: **Fc gamma Receptors: Old Friends and New Family Members.** *Immunity* 2006, **24**:19-28.
35. Lubeck MD, Steplewski Z, Baglia F, Klein MH, Dorrington KJ, Koprowski H: **The interaction of murine IgG subclass proteins with human monocyte Fc receptors.** *J Immunol* 1985, **135**:1299-1304.
36. Lubeck MD, Steplewski Z, Baglia F, Klein MH, Dorrington KJ, Koprowski H: **The interaction of murine IgG subclass proteins with human monocyte Fc receptors.** *J Immunol* 1985, **135**:1299-1304.
37. Jones DH, Looney RJ, Anderson CL: **Two distinct classes of IgG Fc receptors on a human monocyte line (U937) defined by differences in binding of murine IgG subclasses at low ionic strength.** *J Immunol* 1985, **135**:3348-3353.
38. Dougherty GJ, Selvendran Y, Murdoch S, Palmer DG, Hogg N: **The human mononuclear phagocyte high-affinity Fc receptor, FcRI, defined by a monoclonal antibody, 10.1.** *Eur J Immunol* 1987, **17**:1453-1459.
39. Spaggiari GM, Contini P, Carosio R, Arvigo M, Ghio M, Oddone D, Dondero A, Zocchi MR, Puppo F, Indiveri F, Poggi A: **Soluble HLA class I molecules induce natural killer cell apoptosis through the engagement of CD8: evidence for a negative regulation exerted by members of the inhibitory receptor superfamily.** *Blood* 2002, **99**:1706-1714.
40. Chapman TL, Heikema AP, West AP Jr, Bjorkman PJ: **Crystal structure and ligand binding properties of the D1D2 region of the inhibitory receptor LIR-I (ILT2).** *Immunity* 2000, **13**:727-736.
41. Bosselut R, Kubo S, Guintert T, Kopacz JL, Altman JD, Feigenbaum L, Singer A: **Role of CD8beta domains in CD8 coreceptor function: importance for MHC I binding, signaling, and positive selection of CD8+ T cells in the thymus.** *Immunity* 2000:409-418.
42. Daniels MA, Devine L, Miller JD, Moser JM, Lukacher AE, Altman JD, Kavathas P, Hogquist KA, Jameson SC: **CD8 binding to MHC class I molecules is influenced by T cell maturation and glycosylation.** *Immunity* 2001, **15**:1051-1061.
43. Moody AM, Chui D, Reche PA, Priatel JJ, Marth JD, Reinherz EL: **Developmentally regulated glycosylation of the CD8alpha beta coreceptor stalk modulates ligand binding.** *Cell* 2001, **107**:501-512.
44. Maeurer MJ, Necker A, Salter RD, Castelli C, Hohn H, Karbach J, Freitag K, Neukirch C, Knuth A, Jager E: **Improved detection of melanoma antigen-specific T cells expressing low or high levels of CD8 by HLA-A2 tetramers presenting a Melan-A/Mart-1 peptide analogue.** *Int J Cancer* 2002, **97**:64-71.
45. Kuball J, Schmitz FW, Voss RH, Ferreira EA, Engel R, Guillaume P, Strand S, Romero P, Huber C, Sherman LA, Theobald M: **Cooperation of human tumor-reactive CD4+ and CD8+ T cells after redirection of their specificity by a high-affinity p53A2.1-specific TCR.** *Immunity* 2005, **22**:117-129.
46. Campanelli R, Palermo B, Garbelli S, Mantovani S, Lucchi P, Necker A, Lantelme E, Giachino C: **Human CD8 co-receptor is strictly involved in MHC-peptide tetramer-TCR binding and T cell activation.** *Int Immunol* 2002, **14**:39-44.
47. Pittet MJ, Rubio-Godoy V, Bioley G, Guillaume P, Batard P, Speiser D, Luescher I, Cerottini JC, Romero P, Zippelius A: **Alpha 3 domain mutants of peptide/MHC class I multimers allow the selective isolation of high avidity tumor-reactive CD8 T cells.** *J Immunol* 2003, **171**:1844-1849.
48. Snow PM, Van de RM, Terhorst C: **Association between the human thymic differentiation antigens T6 and T8.** *Eur J Immunol* 1985, **15**:529-532.
49. Barber EK, Dasgupta JD, Schlossman SF, Trevillyan JM, Rudd CE: **The CD4 and CD8 antigens are coupled to a protein-tyrosine kinase (p56lck) that phosphorylates the CD3 complex.** *Proc Natl Acad Sci USA* 1989, **86**:3277-3281.
50. Snow PM, Keizer G, Coligan JE, Terhorst C: **Purification and N-terminal amino acid sequence of the human T cell surface antigen T8.** *J Immunol* 1984, **133**:2058-2066.
51. Pichler WJ, Wolff-Vorbeck G, Birke C, Rieber P, Peter HH: **Comparison of the inhibitory activity of seven anti-T8 antibodies on specific cellular cytotoxicity.** *Immunobiology* 1984, **167**:376-388.
52. Biddison WE, Rao PE, Talle MA, Boselli CM, Goldstein G: **Distinct epitopes on the T8 molecule are differentially involved in cytotoxic T cell function.** *Hum Immunol* 1984, **9**:117-130.
53. Van Severter GA, Van Lier RA, Spits H, Ivanyi P, Melief CJ: **Evidence for a regulatory role of the T8 (CD8) antigen in antigen-specific and anti-T3-(CD3)-induced lytic activity of allospecific cytotoxic T lymphocyte clones.** *Eur J Immunol* 1986, **16**:1363-1371.
54. Fridlender ZG, Rabinowitz R, Schlesinger M: **Monocytes confer CD14 antigenicity on activated lymphocytes.** *Hum Immunol* 1999, **60**:1028-1038.
55. Tartakovsky B, Fried M, Bleiberg M, Turner D, Hoffman M, Yust I: **An intracellular antigen that reacts with MO2, a monoclonal antibody to CD14, is expressed by human lymphocytes.** *Immunol Lett* 2003, **85**:35-40.
56. Pascale MC, Malagolini N, Serafini-Cessi F, Migliaccio G, Leone A, Bonatti S: **Biosynthesis and oligosaccharide structure of human CD8 glycoprotein expressed in a rat epithelial cell line.** *J Biol Chem* 1992, **267**:9940-9947.
57. ten Brinke A, Vaandrager AB, Haagsman HP, Ridder AN, van Golde LM, Batenburg JJ: **Structural requirements for palmitoylation of surfactant protein C precursor.** *Biochem J* 2002, **361**:663-671.
58. Wu W, Harley PH, Punt JA, Sharrow SO, Kears KP: **Identification of CD8 as a peanut agglutinin (PNA) receptor molecule on immature thymocytes.** *J Exp Med* 1996, **184**:759-764.
59. Arcaro A, Gregoire C, Bakker TR, Baldi L, Jordan M, Goffin L, Boucheron N, Wurm F, van der Merwe PA, Malissen B, Luescher IF: **CD8beta endows CD8 with efficient coreceptor function by coupling T cell receptor/CD3 to raft-associated CD8/p56(lck) complexes.** *J Exp Med* 2001, **194**:1485-1495.
60. Vremec D, Shortman K: **Dendritic cell subtypes in mouse lymphoid organs: cross-correlation of surface markers, changes with incubation, and differences among thymus, spleen, and lymph nodes.** *J Immunol* 1997:565-573.
61. Vremec D, Zorbas M, Scollay R, Saunders DJ, Ardavin CF, Wu L, Shortman K: **The surface phenotype of dendritic cells purified from mouse thymus and spleen: investigation of the CD8 expression by a subpopulation of dendritic cells.** *J Exp Med* 1992, **176**:47-58.
62. Leishman AJ, Naidenko OV, Attinger A, Koning F, Lena CJ, Xiong Y, Chang HC, Reinherz E, Kronenberg M, Cheroutre H: **T cell responses modulated through interaction between CD8alpha alpha and the nonclassical MHC class I molecule, TL.** *Science* 2001:1936-1939.
63. Gangadharan D, Cheroutre H: **The CD8 isoform CD8alpha alpha is not a functional homologue of the TCR co-receptor CD8alpha beta.** *Curr Opin Immunol* 2004, **16**:264-270.
64. Romero P, Cerottini JC, Luescher I: **On the significance of CD8alpha alpha expression for T cell memory.** *Eur J Immunol* 2005, **35**:3092-3094.
65. Madakamutil LT, Christen U, Lena CJ, Wang-Zhu Y, Attinger A, Sundarrajan M, Ellmeier W, von Herrath MG, Jensen P, Littman DR, Cheroutre H: **CD8alpha alpha-mediated survival and differentiation of CD8 memory T cell precursors.** *Science* 2004, **304**:590-593.
66. Chandele A, Kaech SM: **Cutting edge: memory CD8 T cell maturation occurs independently of CD8alpha alpha.** *J Immunol* 2005, **175**:5619-5623.
67. Zhong W, Reinherz EL: **CD8 alpha alpha homodimer expression and role in CD8 T cell memory generation during influ-**

- enza virus A infection in mice. *Eur J Immunol* 2005, **35**:3103-3110.
68. Leishman AJ, Naidenko OV, Attinger A, Koning F, Lena CJ, Xiong Y, Chang HC, Reinherz E, Kronenberg M, Cheroutre H: **T cell responses modulated through interaction between CD8alpha and the nonclassical MHC class I molecule, TL.** *Science* 2001:1936-1939.
 69. Gabert J, Langlet C, Zamoyska R, Parnes JR, Schmitt-Verhulst AM, Malissen B: **Reconstitution of MHC class I specificity by transfer of the T cell receptor and Lyt-2 genes.** *Cell* 1987, **50**:545-554.
 70. Holler PD, Kranz DM: **Quantitative analysis of the contribution of TCR/pepMHC affinity and CD8 to T cell activation.** *Immunity* 2003, **18**:255-264.
 71. Letourneur F, Gabert J, Cosson P, Blanc D, Davoust J, Malissen B: **A signaling role for the cytoplasmic segment of the CD8 alpha chain detected under limiting stimulatory conditions.** *Proc Natl Acad Sci USA* 1990, **87**:2339-2343.
 72. Wheeler CJ, von Hoegen P, Parnes JR: **An immunological role for the CD8 beta-chain.** *Nature* 1992, **357**:247-249.
 73. Zamoyska R, Derham P, Gorman SD, von Hoegen P, Bolen JB, Veillette A, Parnes JR: **Inability of CD8 alpha' polypeptides to associate with p56lck correlates with impaired function in vitro and lack of expression in vivo.** *Nature* 1989, **342**:278-281.
 74. Chu DH, Spits H, Peyron JF, Rowley RB, Bolen JB, Weiss A: **The Syk protein tyrosine kinase can function independently of CD45 or Lck in T cell antigen receptor signaling.** *EMBO J* 1996, **15**:6251-6261.
 75. Taylor N, Jahn T, Smith S, Lamkin T, Uribe L, Liu Y, Durden DL, Weinberg K: **Differential activation of the tyrosine kinases ZAP-70 and Syk after Fc gamma RI stimulation.** *Blood* 1997, **89**:388-396.
 76. Tridandapani S, Lyden TW, Smith JL, Carter JE, Coggeshall KM, Anderson CL: **The adapter protein LAT enhances fc gamma receptor-mediated signal transduction in myeloid cells.** *J Biol Chem* 2000, **275**:20480-20487.
 77. Kuster H, Thompson H, Kinet JP: **Characterization and expression of the gene for the human Fc receptor gamma subunit. Definition of a new gene family.** *J Biol Chem* 1990, **265**:6448-6452.
 78. Saitoh S, Odom S, Gomez G, Sommers CL, Young HA, Rivera J, Samelson LE: **The four distal tyrosines are required for LAT-dependent signaling in Fc epsilon RI-mediated mast cell activation.** *J Exp Med* 2003, **198**:831-843.
 79. Barrett TA, Gajewski TF, Danielpour D, Chang EB, Beagley KW, Bluestone JA: **Differential function of intestinal intraepithelial lymphocyte subsets.** *J Immunol* 1992, **149**:1124-1130.
 80. Hirji N, Lin TJ, Bissonnette E, Belosevic M, Befus AD: **Mechanisms of macrophage stimulation through CD8: macrophage CD8alpha and CD8beta induce nitric oxide production and associated killing of the parasite Leishmania major.** *J Immunol* 1998, **160**:6004-6011.
 81. Lin TJ, Hirji N, Stenton GR, Gilchrist M, Grill BJ, Schreiber AD, Befus AD: **Activation of macrophage CD8: pharmacological studies of TNF and IL-1 beta production.** *J Immunol* 2000, **164**:1783-1792.
 82. Tam FW, Smith J, Morel D, Karkar AM, Thompson EM, Cook HT, Pusey CD: **Development of scarring and renal failure in a rat model of crescentic glomerulonephritis.** *Nephrol Dial Transplant* 1999, **14**:1658-1666.
 83. May E, Dorris ML, Satumtira N, Iqbal I, Rehman MI, Lightfoot E, Taugrog JD: **CD8 alpha beta T cells are not essential to the pathogenesis of arthritis or colitis in HLA-B27 transgenic rats.** *J Immunol* 2003, **170**:1099-1105.
 84. Bonnotte B, Larmonier N, Favre N, Fromentin A, Moutet M, Martin M, Gurbuxani S, Solary E, Chauffert B, Martin F: **Identification of tumor-infiltrating macrophages as the killers of tumor cells after immunization in a rat model system.** *J Immunol* 2001, **167**:5077-5083.
 85. Schroeter M, Stoll G, Weissert R, Hartung HP, Lassmann H, Jander S: **CD8+ phagocyte recruitment in rat experimental autoimmune encephalomyelitis: association with inflammatory tissue destruction.** *Am J Pathol* 2003, **163**:1517-1524.
 86. Schroeter M, Jander S, Huitinga I, Stoll G: **CD8+ phagocytes in focal ischemia of the rat brain: predominant origin from hematogenous macrophages and targeting to areas of pan-necrosis.** *Acta Neuropathol (Berl)* 2001, **101**:440-448.
 87. Karkar AM, Smith J, Pusey CD: **Prevention and treatment of experimental crescentic glomerulonephritis by blocking tumour necrosis factor-alpha.** *Nephrol Dial Transplant* 2001, **16**:518-524.
 88. Braun J, Brandt J, Listing J, Zink A, Alten R, Golder W, Gromnica-Ihle E, Kellner H, Krause A, Schneider M, Sorensen H, Zeidler H, Thriene W, Sieper J: **Treatment of active ankylosing spondylitis with infliximab: a randomised controlled multicentre trial.** *Lancet* 2002, **359**:1187-1193.
 89. Paulnock DM: **Macrophages: Practical Approach.** Oxford: Oxford University Press; 2000.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

